

Wisdom and Influence of Elders: Possibilities for Health Promotion and Decreasing Tobacco Exposure in First Nations Communities

Colleen Varcoe, PhD, RN,¹ Joan L. Bottorff, PhD, RN, FCAHS,² Joanne Carey, MA,³ Debbie Sullivan, BSN, MN,⁴ Wanda Williams⁵

ABSTRACT

Objectives: This community-based project examined the interpersonal and system influences on smoking practices and exposure to cigarette smoke that place young pregnant and parenting First Nations women and their children at risk. The aim was to work with the community using this information to develop meaningful strategies to support tobacco reduction and reduce exposure to second-hand smoke.

Method: Ethnographic methods informed by decolonizing approaches were used to study smoking practices and exposure within the context of six First Nations rural reserve villages in the Gitx̱san territory in North-Western British Columbia. Data were collected through individual interviews with key community members, individual and group interviews with women, men, youth and older people, including Elders, and participant observation.

Findings: Older people, including grandparents and Elders expressed great interest in being more involved in their communities, particularly with youth. Other community members, especially mothers, saw such older people as important in diminishing the tobacco exposure problem, with ideas for their involvement ranging from role modeling non-smoking or respectful smoking to teaching cultural traditions.

Conclusion: Elders in First Nations communities represent a culturally-relevant resource for health that is currently underused. Communities might draw upon their wisdom and influence to decrease tobacco exposure and promote community health in other ways. This offers a concrete, practical strategy in response to widespread calls for supporting the culture of Aboriginal people as part of health promotion.

Key words: Indigenous population; tobacco; passive smoking; mothers; pregnant women; health promotion

La traduction du résumé se trouve à la fin de l'article.

Can J Public Health 2010;101(2):154-58.

High rates of maternal smoking during pregnancy and postpartum and exposure of infants and children to second-hand smoke (SHS) are persistent health concerns that contribute to health disparities between Aboriginal (encompassing First Nations, Métis and Inuit people)¹ and non-Aboriginal Canadians.^{2,3} Aboriginal women are about three times more likely to be smoking during pregnancy as non-Aboriginal women.^{4,5} Aboriginal households report that 32% (compared to provincial rates of 18%) of households with children under age 11 experience daily or nearly daily exposure to SHS.⁶ Similar to indigenous populations worldwide, rates of cigarette smoking and exposure to SHS in Canadian Aboriginal communities are influenced by levels of poverty and community resources, geographic isolation, and historical and ongoing colonization. In spite of the above, there have been few studies of antenatal smoking and indigenous women.⁷ Interventions aimed at supporting behavioural change for individuals, such as drug therapy and quit-lines, have been studied in Canadian Aboriginal contexts, but shown to have limited impact,⁸⁻¹¹ and recent legal and taxation policies to curb tobacco sales have had limited success.¹²⁻¹⁴ There has been little study to develop context-specific public health interventions. Contemporary Aboriginal communities are actively seeking health and revitalization, in part through emphasizing culture and tradition.^{15,16} Members of the Gitx̱san First Nations were interested in finding ways to protect pregnant women and those with young children from cigarette smoke and to support women's tobacco reduction efforts. This participatory ethnographic study examined the research questions in Table 1; this paper reports on one context-specific strategy.

METHOD

Using an ethnographic approach within a participatory paradigm informed by decolonizing intent,^{17,18} community health care providers partnered with academic researchers to explore interpersonal and system influences on smoking practices and exposure to SHS that increase health risk for pregnant and parenting women and their children and to develop meaningful strategies to support tobacco reduction and reduce SHS exposure.¹⁹ Drawing on principles for conducting research with Aboriginal communities,^{20,21} ethical approval was obtained from the community and the researchers' university.

Context

The six study communities were in Gitx̱san Territory in rural northern British Columbia (Table 2). Cigarette smoking was prevalent, particularly among youth.^{22,23} Economic factors of the reserve system coupled with limited employment income and poverty made

Author Affiliations

1. Associate Professor, School of Nursing, University of British Columbia, Vancouver, BC
2. Professor, Institute for Healthy Living and Chronic Disease Prevention, University of British Columbia Okanagan, Kelowna, BC
3. Research Coordinator, Institute for Healthy Living and Chronic Disease Prevention, University of British Columbia Okanagan, Kelowna, BC
4. Community Health Nurse, Gitsegukla Health Programs and Services, Gitsegukla, BC
5. Community Health Representative, Gitsegukla Health Programs and Services, Gitsegukla, BC

Correspondence: Dr. Colleen Varcoe, Associate Director, Research, University of British Columbia School of Nursing, T201-2211 Wesbrook Mall, Vancouver, BC V6T 2B5, Tel: 604-827-3121, Fax: 604-822-7466, E-mail: colleen.varcoe@nursing.ubc.ca

Conflict of Interest: None to declare.

Table 1. Research Questions

1. How does the social and physical context (i.e., gender, age, culture, living on reserve, rural/urban) influence young pregnant and parenting Aboriginal women's tobacco use, and young women's and children's exposure to SHS by other smokers?
2. What strategies do young Aboriginal women use to reduce tobacco use and to minimize SHS exposure for themselves and their children?
3. What factors and processes need to be considered in designing interventions, programs and policies to support smoke-free spaces and tobacco reduction that will benefit young pregnant and parenting Aboriginal women?
4. What are community reactions to knowledge gained about the experiences of young pregnant and parenting women related to tobacco, and identified strategies and priorities for tobacco reduction?
5. What are the challenges and opportunities for providing smoke-free environments for young pregnant and parenting Aboriginal women and their children as viewed by community members?
6. How can the research process and the findings inform the development of context-specific strategies and action plans for tobacco reduction that address the needs of young pregnant and parenting Aboriginal women and their children?

tobacco both a source of income and a reasonably affordable source of pleasure. Further, financial factors and limited affordable recreational activities contributed to bingo being a popular social and fundraising activity that involved smoking and exposure to SHS in all but one community. Community leadership was diverse, encompassing band councils, elected chiefs and hereditary chiefs who operated through different mechanisms of influence, facing competing priorities that were not necessarily focused on health issues such as tobacco.

Data collection

Multiple participatory strategies were used (Table 3). Data were collected through individual and focus group interviews and participant observation of activities including community walks, prenatal programs, parenting and lunch programs. Participants were recruited using purposive sampling to optimize diversity. As shown in Table 4, the 66 participants included 26 young women who were pregnant or parenting young children, 9 Elders, 6 youth, 10 adult community members and 15 key community members (KCM), all of whom had prominent roles and some of whom also were considered Elders by the community. Interviews and focus groups were conducted by a trained research assistant who was a young woman with children and a member of the Gitx̄san Nation; about half of the sessions were co-conducted with other team members. All interviews were recorded and transcribed. Team members conducted participant observations and kept detailed field notes.

Data analysis

Data were analyzed using ethnographic techniques.^{24,25} Team members read interviews repeatedly and identified and categorized meaning units. Categories were compared and contrasted to develop themes. Based on several initial interviews, an inductive coding scheme was developed through consensus. Interview data from diverse participants across all study communities were compared and field note data supplemented interview data to develop themes and descriptions. Investigator subgroups explored particular themes in greater depth. Procedures were documented to develop an audit trail regarding how conclusions were drawn. Community-based research team members participated in all aspects of analysis, and community members provided feedback through an interactive event that used the findings.

Table 2. Community Profile³¹

Population	Study villages range from 480 to 2000
Distances	75 km between the largest village and the closest urban centre with airport, shopping, hospital, recreation facilities (population approximately 5,000) Study villages are clustered within a 60 km radius.
Income and Employment	Average annual income Women: \$9100-\$14,688 Men: \$8144-\$10,816 Unemployment rate 30-50%

Table 3. Participatory and Context-appropriate Strategies

Phase	Strategy
Recruitment	<ul style="list-style-type: none"> • Community-based team members (community health nurse [CHN], community health representative [CHR], research assistant) recruited in multiple community venues (e.g., health fairs, job fairs) • Family members welcomed to participate together (e.g., cousins, mother and daughters, sisters)
Interviewing	<ul style="list-style-type: none"> • Community-based First Nations team members present at all interviews with First Nations members • Series of small group interviews held with young women to enhance comfort • Food routinely provided • Gitx̄san-speaking team member present for Elders focus group (interpreting intermittently)
Analysis	<ul style="list-style-type: none"> • Analysis conducted throughout data collection • All team members (community-based and academic) engaged throughout (from initial analysis to writing of papers) • Multiple methods used to take developing analysis to community (regular newsletters, video, community events) for feedback
Knowledge Exchange	<ul style="list-style-type: none"> • Participatory community event using actors and video • Regular newsletters

FINDINGS

Overall findings of this study highlighted how possibilities for tobacco reduction must be understood within the historical and socio-cultural context of tobacco use in rural First Nations communities and drew attention to context-specific strategies.¹⁹ One such strategy identified throughout the study was to engage Elders. Although cultural erosion has affected the status of Elders, the latter continue to play dynamic and influential roles. Although some Elders smoked, and some expressed doubt about their potential for influence, success stories regarding tobacco control were routinely linked to Elders, and community members expressed hope that Elders could provide further leadership.

The Role of Elders

In the Gitx̄san tradition, Elders are people knowledgeable about culture and tradition who can articulate “the way” – beliefs and expectations of the Gitx̄san people. For example, Elders would know the Feast system and the relations among various clans. Elders are not always persons over a certain age, and not all older persons are considered Elders. Rather, Elders are those who have and show concern for others and the community and show leadership. The position of Elder in the Gitx̄san nation is one of esteem and respect. However, the position is also influenced by politics. That is, certain Elders are more likely to be called upon than others because of their connections, and some people assert themselves as Elders rather than achieving that status through community recognition.

In the study communities, Elders were frequently invited to speak to youth in classrooms and share stories with children so as to maintain cultural knowledge. Importantly, participants thought Elders had a key role in guiding youth:

Table 4. Participant Demographics*

	n	Participation	Age	Smoking Status
Young women (all parenting; 3 pregnant at time of interview)	26	2-4 individual and small group interviews	17-34 (m†=24.8)	Ex-smoker (n=7) Occasional (n=9) Daily (n=10)
Key community members (KCM) (e.g., Public/Community health nurses, community health representatives [CHR], family facilitators, addiction counsellors, educators, managers)	15 (13 female; 2 male)	Individual interview	32-64 (m=51)	Never smoked (n=7) Ex-smoker (n=2) Occasional (n=5) Daily (n=1)
Elders	9 (8 female; 1 male)	Focus group interview	63-87 (m=75)	Never smoked (n=4) Ex-smoker (n=2) Occasional (n=2) Daily (n=1)
Community members (recruited to increase representation from all communities)	10 (7 female; 3 male)	Focus group interview	27-61 (m=43.5)	Ex-smoker (n=3) Occasional (n=3) Daily (n=4)
Youth	6 (3 female; 3 male)	Focus group interview	13-17 (m=15.3)	Never smoked (n=4) Ex-smoker (n=2)

* all participants were members of Gitksan First Nation except 4 “Key Community Members”

† m=mean

I tell children... if an Elder comes and talks to you about some things you're doing, it's more than likely they're giving you a detour of your life you know, they're saying I've been down that road, don't bother with it, try something else because you're just wasting time and energy and probably sacrificing your health, your youth especially. [KCM]

In interviews, Elders expressed great interest in being more involved in their communities, particularly with youth. One said, “I would be interested in being available...for example...if there were teenage moms that just needed someone that they could talk to or spend time with.” However, alongside the professed reverence for Elders, many participants, including Elders themselves, discussed the erosion of the role of Elders and declining respect.

I was taught to always respect Elders. And then when I see younger people totally disrespecting them in their tone of voice or just the way they speak to them, I just get really angry and frustrated inside and go, “didn't your parents teach you?!” [Young mother]

Despite these tensions, participants routinely identified Elders as an important possible source of influence.

Contemporary success stories linked to role of Elders

Enduring respect and concern for Elders and their instrumental role in positive tobacco control measures were evident in the data. For example, a locally produced anti-smoking poster was well known for the Elder's message: “We smoke fish, not tobacco.” In one community, reports about efforts to establish smoke-free bingos in the community hall were directly linked to the support of Elders.

Well, I was surprised that it was an Elders' group that brought up that the bingos should be smoke-free. Like I thought I would have heard that from some moms, but I think I first heard it through an Elders' discussion. [Young mother]

In another community, a chief recounted banning smoking at feasts to protect the health of Elders:

When I became a chief and we were going to have a feast... someone said to me, “gee, you know I don't really mind coming to feasts, but it's the smoke I can't stand,” you know. So, there was an opportunity for me because I took the name, and I stood up there, and I said, “this is going to be a non-smoking feast because there are Elders that are here and they enjoy the companionship of everybody, but there's a big difficulty in smoking and most of our Elders are sick...so no more smoking in our feast.” So, there's been no smoking in most of all our feasts....since '95, I think, when I took the name. [Elder]

In addition to successes at a collective or policy level, participants told stories of Elders being inspirational. One participant told how an Elder had thrown his cigarettes into the ocean, quitting smoking for good: “for kids to hear that, I think ... can be very inspiring to them, you know.”

Elders as potential source of leadership

In discussing strategies for extending smoking restrictions to other settings and events, the importance of enlisting the support of Elders was a consistent theme. Community members, especially mothers, saw Elders as important allies in dealing with SHS. Ideas for their involvement ranged from role modeling non-smoking or respectful smoking to teaching broader traditions. With limited housing options, multiple generations often lived together, meaning that Elders had potential for influence directly in the home. Many noted that Elders with first-hand experiences of the harmful effects of smoking had made important changes that served as a model:

Most of the Elders that I talk to don't smoke or don't like it. They smoked so much in their lives that they just quit 'cause they're Elders now, and it's starting to affect their health. [Young mother]

Some Elders had stopped or reduced smoking for their grandchildren. Of those who continued to smoke, some were adamant that youth should not take up the habit and worried about children being exposed to SHS.

Although many talked about the importance of smoke-free community events to protect the failing health of Elders, those who wanted to extend smoking restrictions wanted Elders to do more.

I think once our Elders step in and show what needs to be done and stuff like, like even though our Elders don't smoke, but if they were to enforce no smoking here, like even right out of the bingo halls you know. It would step up and show that our Elders are serious... [Young mother]

Participants thought that drawing on Elders as a source of leadership in relation to smoking and exposure to SHS would not only be culturally appropriate, but also would be congruent with how the community understood the causes of smoking including loss of traditions, underemployment and boredom. One KCM emphasized the importance of such indirect measures and actively engaging the community, saying, “because then you're starting to look at under-

pimming issues and root causes rather than [something] as specific as changing eating or changing behaviour around smoking."

Since the completion of data collection, one of the largest bingo halls, attached to a school, was pressured by community action to become smoke-free. Community members noted that the protest that led to the change was initiated by a young mother and supported by community Elders. The women saw this change as significant, not only as symbolic of rising concern and action, but as affecting one of their limited social opportunities that in turn influenced their ability to reduce their own smoking.¹⁹

Study limitations

Conducting the study with one specific First Nation limits the applicability of the findings to other groups. The focus on tobacco reduction may have constrained participants who smoked from expressing certain viewpoints; however, collecting data in both individual and focus group formats and from people with a range of smoking experiences, increased data variation. Although generalizability is inappropriate given the sample size and specific context, the fit with other literature suggests these findings have applicability beyond the Gitksan, particularly to other First Nations and to other health concerns beyond tobacco.

DISCUSSION

Although First Nations communities are diverse, representing over 50 different language groups and diverse multiple cultures,^{26,27} many communities share a high value for Elders and see them as spiritual leaders. First Nations also share the experiences of colonization – and while those experiences vary, all Aboriginal people were subject to colonial policies flowing from the Indian Act that eroded their cultures, languages, ways of life and access to material resources. Addressing these dynamics that have led to poor health for many people must be central to health promotion. Other researchers have argued that strengthening ethnocultural identity, community integration and political empowerment can contribute to improving mental health for Aboriginal people.²⁸⁻³⁰ We suggest that these have broad health impacts and that engaging Elders is one key element in such strengthening. Researchers need to recognize Elders and their importance in conducting research – and a similar approach needs to be taken to tobacco control.

As interventions aimed at behavioural change have had limited impact on tobacco control, new models are required. Our findings indicate that for First Nations communities, drawing on the Elders may be a key strategy for such models. Rather than beginning with predetermined tobacco reduction strategies from other contexts, Elders may guide context-specific approaches, including using their own influence through multiple connections. Importantly, drawing on the wisdom of Elders contributes to cultural strength and therefore would be an approach that would help address what the participants saw as the root cause of smoking: cultural erosion.

Appropriate protocols and approaches to involving Elders will vary with each community and nation. Invitations to Elders to participate in health strategies such as tobacco reduction measures need to be made in the context of meaningful relationships. Local knowledge would be required both to know who to approach and how, and to build bridges to establish strong relationships with Elders. Programs to partner youth and Elders – helping Elders with firewood was suggested – could have multiple benefits. Understanding what

constitutes an Elder and identifying which people are considered Elders by whom are prerequisite to seeking consultation. Existing community health advisories could be used to expand or formalize the role of Elders. Further, broad and multiple strategies are needed in order to involve Elders beyond those who are routinely identified to give input. In some contexts, consultation with as many Elders as possible might be appropriate; in others, a snowball approach to seek out diverse Elders might be best. Our experience suggests that concrete strategies such as insuring that bilingual community members are available to interpret, inviting family members together, and offering door-to-door transportation are essential.

Realistically enacting such an orientation to health promotion would require commitment at all levels. First, funding bodies such as Health Authorities could require guidance from Elders in relation to health programs, including those aimed at tobacco reduction. Bodies funding health research could specify similar requirements. Second, researchers and people implementing programs routinely could build in provisions for drawing on such guidance. In settings such as universities, colleges and some health care settings, Resident Elders are becoming increasingly formalized and paid positions, and such strategies might be adapted for community organizations, particularly if support is provided through program and research funding. Finally, service providers and community members could identify Elders who can provide wisdom and leadership.

REFERENCES

1. Royal Commission on Aboriginal Peoples. *Report of the Royal Commission on Aboriginal Peoples*. Ottawa, ON: Canada Communications Group-Publishing, 1996.
2. Heaman MI, Chalmers K. Prevalence and correlates of smoking during pregnancy: A comparison of Aboriginal and Non-Aboriginal women in Manitoba. *Birth* 2005;32(4):299-305.
3. Wenman WM, Joffres MR, Tataryn IV. A prospective cohort study of pregnancy risk factors and birth outcomes in Aboriginal women. *CMAJ* 2004;171(6):585-89.
4. Muhajarine N, D'Arcy C, Edouard L. Prevalence and predictors of health risk behaviours during early pregnancy: Saskatoon Pregnancy and Health Study. *Can J Public Health* 1997;88(6):375-79.
5. Heaman MI, Chalmers K. Prevalence and correlates of smoking during pregnancy: A comparison of Aboriginal and non-Aboriginal women in Manitoba. *Birth: Issues in Perinatal Care* 2005;32(4):299-305.
6. Angus Reid Group. *Tobacco Use in BC*. Vancouver, BC: Heart and Stroke Foundation of BC & Yukon, 1997.
7. Gilligan C, Sanson-Fisher R, Eades S, D'Este C. Antenatal smoking in vulnerable population groups: An area of need. *J Obstet Gynaecol* 2007;27:664-71.
8. Wardman AED, Khan NA. Tobacco cessation pharmacotherapy use among First Nations persons residing within British Columbia. *Nicotine Tob Res* 2004;6(4):689-92.
9. Wardman D, Quantz D, Tootoosis J, Khan N. Tobacco cessation drug therapy among Canada's Aboriginal people. *Nicotine Tob Res* 2007;9(5):607-11.
10. Donelle L, Hoffman-Goetz L. An exploratory study of Canadian Aboriginal online health care forums. *Health Commun* 2008;23(3):270-81.
11. Hayward LM, Campbell HS, Sutherland-Brown C. Aboriginal users of Canadian quilines: An exploratory analysis. *Tobacco Control* 2007;16:i1-i2.
12. Fiddler C. Feds enforce new tobacco strategy which targets Aboriginal communities. *Windspeaker* 2008;26(3):8-9.
13. Luk R, Cohen JE, Ferrence R, McDonald PW, Schwartz R, Bondy SJ. Prevalence and correlates of purchasing contraband cigarettes on First Nations reserves in Ontario, Canada. *Addiction* 2009;104(3):488-95.
14. Kondro W. Coalition urges crackdown on contraband cigarettes. *CMAJ* 2007;176(11):1569.
15. Baker A, Ivers RG, Bowman J, Butler T, Kay-Lambkin FJ, Wye P, et al. Where there's smoke, there's fire: High prevalence of smoking among some sub-populations and recommendations for intervention. *Drug & Alcohol Review* 2006;25:85-96.
16. McCabe G. Mind, body, emotions and spirit: Reaching to the ancestors for healing. *Counseling Psychol Q* 2008;21(2):143-52.
17. Smith LT. *Decolonizing methodologies: Research and indigenous people*. Dunedin: University of Otago Press, 1999.

18. Denzin NK, Lincoln YS, Smith LT (Eds.). *Handbook of Critical and Indigenous Methodologies*. Thousand Oaks, CA: Sage, 2008.
19. Bottorff JL, Carey J, Mowatt R, Varcoe C, Johnson JL, Hutchinson P, et al. Bingo halls and smoking: Perspectives of First Nations women. *Health & Place* 2009;15:1014-21.
20. Interagency Advisory Panel on Research Ethics. Draft 2nd Edition of the Tri-council Policy Statement: Ethical Conduct for Research Involving Humans. Ottawa, ON: Interagency Secretariat on Research Ethics, 2008.
21. Canadian Institutes of Health Research. CIHR Guidelines for Health Research Involving Aboriginal People. 2008.
22. Ho K, Blanchard C. British Columbia Youth Survey on Smoking & Health II: The Results (A Report for Teens by Teens). Vancouver, BC: School of Nursing, University of British Columbia, 2004.
23. Johnson JL, Tucker R, Ratner PA, Bottorff JL, Prkachin KM, Shoveller J, Zumbo B. Socio-demographic correlates of cigarette smoking among high school students: Results from the British Columbia Youth Survey on Smoking and Health. *Can J Public Health* 2004;95:268-71.
24. Atkinson P, Hammersley M. Ethnography and participant observation. In: Denzin NK, Lincoln YS (Eds.), *Handbook of Qualitative Research*. Thousand Oaks, CA: Sage, 1994;248-61.
25. Hammersley M, Atkinson P. *Ethnography: Principles in Practice*, 2nd ed. London: Routledge, 1995.
26. Statistics Canada. Aboriginal peoples in Canada in 2006: Inuit, Métis and First Nations, 2006 Census. Ottawa, ON: Statistics Canada, 2008.
27. Cook E-D, Howe D. Aboriginal Languages of Canada. In: O'Grady W, Archibald J (Eds.), *Contemporary Linguistic Analysis*, 5th ed. Toronto, ON: Addison Wesley Longman, 2004;294-309.
28. Richmond CAM, Ross NA. The determinants of First Nation and Inuit health: A critical population health approach. *Health & Place* 2009;15(2):403-11.
29. Kirmayer L, Simpson C, Cargo M. Indigenous populations healing traditions: Culture, community and mental health promotion with Canadian Aboriginal peoples. *Australasian Psychiatry* 2003;11:515.
30. Chandler MJ, Lalonde C. Cultural continuity as a hedge against suicide in Canada's First Nations. *Transcultural Psychiatry* 1998;35(2):191-219.
31. Statistics Canada. Community Profiles. 2006. Available at: <http://www12.statcan.ca/english/census06/data/profiles/community/Index> (Accessed October 6, 2008).

Received: October 14, 2009
 Accepted: January 27, 2010

RÉSUMÉ

Objectif : Ce projet communautaire portait sur les influences interpersonnelles et systémiques qui exposent les jeunes femmes enceintes et les jeunes mères des Premières nations et leurs enfants au tabagisme et à la fumée de cigarette. Le but était d'utiliser ces informations en élaborant, avec la collaboration de la communauté, des stratégies concrètes pour réduire le tabagisme et l'exposition à la fumée secondaire.

Méthode : Nous avons employé des méthodes ethnographiques étayées par des approches de décolonisation pour étudier le tabagisme et l'exposition à la fumée de cigarette dans six villages situés dans des réserves rurales des Premières nations sur le territoire gitksan (Nord-Ouest de la Colombie-Britannique). Les données ont été recueillies au moyen d'entretiens individuels avec des membres influents des communautés, d'entretiens individuels et collectifs avec des femmes, des hommes, des jeunes et des personnes âgées (y compris des anciens) et par l'observation des participants.

Résultats : Les personnes âgées, dont les grands-parents et les anciens, étaient vivement intéressées à s'impliquer davantage dans leur communauté, particulièrement auprès des jeunes. Les autres résidents, surtout les mères, considéraient les personnes âgées comme étant importantes pour réduire le problème d'exposition au tabac et avaient des idées pour les impliquer (en étant des exemples de non-fumeurs pour les jeunes ou en leur enseignant le respect du tabac ou d'autres traditions culturelles).

Conclusion : Les anciens des communautés des Premières nations sont une ressource de santé culturellement significative encore sous-utilisée. Les communautés pourraient faire appel à leur sagesse et à leur influence pour réduire l'exposition au tabac et promouvoir la santé communautaire de diverses façons. Ce serait une stratégie concrète et pratique pour répondre aux appels généralisés en faveur de l'intégration de la culture des peuples autochtones dans la promotion de la santé.

Mots clés : population d'origine amérindienne; tabac; tabagisme passif; mères; femmes enceintes; promotion de la santé

**CIHR-IPPH-CPHA
 Call for Population and Public
 Health Research Milestones**

On the occasion of its 100th anniversary, the Canadian Public Health Association (CPHA) has identified 12 public health achievements in an effort to celebrate the contributions of public health from a Canadian perspective and to make our public health history more visible. For more information, please see the CPHA Centenary website (www.cpha100.ca).

To complement this effort, the CIHR-Institute of Population and Public Health (CIHR-IPPH) is partnering with CPHA to initiate a call for milestones in public health research that have significantly contributed to the public's health in Canada and globally.

These milestones in research will be published on the CPHA and IPPH websites and in the IPPH newsletter, and will be featured during CPHA's anniversary month (September 2010) and in a special insert in the *Canadian Journal of Public Health*.

Please visit our website for more information:
<http://www.cihr-irsc.gc.ca/e/41357.html>

**IRSC-ISPP-ACSP Appel de demandes concernant
 des événements marquants liés à la recherche
 en santé publique et des populations**

À l'occasion de son 100^e anniversaire, l'Association canadienne de santé publique (ACSP) a sélectionné 12 réalisations en vue de célébrer les contributions du secteur de la santé publique dans le contexte canadien ainsi que de faire connaître davantage notre histoire dans ce domaine. Pour en savoir plus à ce sujet, consultez le site Web du centenaire de l'ACSP (www.acsp100.ca).

Pour appuyer cet effort, l'Institut de la santé publique et des populations des IRSC (ISPP des IRSC) et l'ACSP lancent un appel conjoint de demandes concernant des événements marquants liés à la recherche en santé publique, événements ayant contribué de manière importante à l'avancement du domaine de la santé publique tant au Canada que sur la scène internationale.

Ces événements seront affichés sur les sites Web de l'ACSP et de l'ISPP. Ils seront également publiés dans le bulletin de l'ISPP et présentés pendant le mois anniversaire de l'ACSP (septembre 2010) et dans un encart spécial de la Revue canadienne de santé publique.

Pour de plus amples renseignements, veuillez visitez notre site Web : <http://www.cihr-irsc.gc.ca/f/41357.html>